

To appear in *The Astrophysical Journal*

Ly α /H α Ratio of Singly Ionized Helium in Quasars¹

Wei Zheng

Center for Astrophysical Sciences, The Johns Hopkins University, Baltimore, MD 21218
zheng@pha.jhu.edu

and

Li-Zhi Fang

Physics Department and Steward Observatory, University of Arizona, Tucson, AZ 85721
fanglz@time.physics.arizona.edu

ABSTRACT

He II Ly α λ 304/H α λ 1640 emission lines are mainly produced by recombination, and their canonical ratio of ~ 10 may be a sensitive reddening indicator. We obtain the high S/N optical spectra of two quasars and combine them with the far-UV spectra that show the He II λ 304 emission. For HS 1700+64, the He II λ 1640 emission is not detected, and an upper limit to it sets the ratio greater than 20. This may not be inconsistent with the theoretical value when all observational uncertainties are taken into consideration. For Q0302-003, the ratio is very low, on the order of unity. The most plausible cause for such a low ratio is extinction in the EUV band by very fine grains of dust. Q0302-003 has a prominent narrow component of FWHM ~ 2000 km s⁻¹ in its major emission lines, and it appears that reddening is associated only with the line-emitting region. We suggest that the geometry of the line-emitting region in high- z quasars resembles that in the low-luminosity active galaxies, with the presence of dust mostly in the outer part.

Subject headings: dust, extinction — quasars; emission lines

1. Introduction

The first UV spectroscopic observation of a quasar (3C 273, Davidsen, Hartig & Fastie 1977) enabled a measurement of the Ly α /H α ratio in an active galactic nucleus (AGN). The low ratio

¹Based on observations with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities of Research in Astronomy, Inc., under NASA contract NAS5-26555.

of ~ 1 was in line with an independent study using the composite spectrum that was derived from various quasars (Baldwin 1977), but was in sharp conflict with a canonical ratio of ~ 10 predicted by standard photoionization models (Osterbrock 1989). This “Ly α /H α problem” raised serious concerns about the validity of photoionization as the main line-emission mechanism in AGN and prompted extensive theoretical interest in the following years. Improved photoionization models (Ferland & Shields 1985, and references therein) invoke large column densities and moderate degrees of ionization. In a partially ionized zone, the low escape probability for Ly α photons makes a high population of excited states, and collisional excitation from these levels enhances Balmer lines. Calculations using a reasonable AGN continuum lead to an enhancement of Balmer lines by a factor of ~ 2 , and hence may not fully explain the observed low Ly α /H α ratio. Another explanation introduces intrinsic reddening in the line-emitting region (MacAlpine 1985). The wavelength-dependent extinction reduces the intensities of observed UV lines, thus lowering the Ly α /H α ratio. The observed Pa α /H α ratio, however, appears to be too low for a straightforward full account as a reddening effect (Puetter et al. 1978). Significant evidence exists for dust in the narrow-line region of Seyfert galaxies (Osterbrock 1989; Netzer & Laor 1993) and for a decrease in extinction with increasing luminosity toward quasars (Cheng, Danese & de Zotti 1983; Rudy et al. 1983). However, Wills et al. (1993) suggested that in intermediate-redshift quasars there may be significant reddening in the narrow-line region.

Accurate assessments of the reddening effect depend on the use of good line pairs whose intrinsic ratios are fairly stable. Since singly ionized helium is hydrogenic, the He II I(Ly α λ 304)/I(H α λ 1640) ratio should therefore be the same as that for hydrogen. The He II emission is produced mainly by recombination, because its excitation level of 40 eV is considerably higher than the average thermal energy in the line-emitting region. The wavelength of He II λ 304 emission coincides with that of O $^{++}$ transition $2p^2\ ^3P_2 - 2p\ 3d\ ^3P_2^0$, allowing Bowen fluorescence radiation (Eastman & MacAlpine 1985; Netzer, Elitzur & Ferland 1985). This radiation mechanism, however, does not appreciably affect the He II ratio itself. The He II λ 304 emission should be extremely sensitive to reddening. While there are no hurdles in the theoretical aspects, it has taken some 20 years to advance from measuring this ratio in hydrogen to that in helium.

2. Observations

As a result of searches for the He II Gunn-Peterson effect in the intergalactic medium, the UV spectra of several hundred quasars have been studied with *HST*. The UV spectra of most high- z quasars are cut off by intervening Lyman-limit systems, and only four quasars have so far been found to exhibit a detectable flux below the wavelength of the redshifted He II λ 304 emission: Q0302-003 ($z=3.286$; Jakobsen et al. 1994), PKS 1935-692 ($z=3.185$; Tytler et al. 1995), HS 1700+64 ($z=2.743$; Davidsen, Kriss & Zheng 1996), and HE 2347-4342 ($z=2.885$; Reimers et al. 1997). The He II λ 304 emission, even with a profile that is partially absorbed, makes it possible to carry out the “Ly α /H α ” test for ionized helium. We used the UV data of HS 1700+64 obtained with the Hopkins

Ultraviolet Telescope (HUT) during the Astro-2 mission and that of Q0302-003 obtained with *HST* Goddard High-Resolution Spectrograph (GHRS) at an improved S/N ratio and resolution (Hogan, Anderson & Rugers 1997). The GHRS data were retrieved from the *HST* archive and co-added.

We obtained the optical spectra of these two quasars on several occasions. The results presented here are mainly from two observing runs in 1996 July and November. At the 2.3m Bok telescope of Steward Observatory, University of Arizona, we used the B&C spectrograph with a Loral 1200×800 CCD detector and a long slit of $2''.5$. On July 16 (UT), the observations of HS 1700+64 were made with a grating of 600 line mm^{-1} for 10,060 s, yielding a resolution of $\sim 3.7 \text{ \AA}$ between ~ 5400 and $7600 \text{ \AA}$. On November 4 and 6 we observed Q0302-003. On the first night, a total exposure of 21,600 s was made using a grating of 832 line mm^{-1} that allows a spectral coverage from ~ 5900 to $7600 \text{ \AA}$ with a resolution of $2.4 \text{ \AA}$. On the other night, the total exposure was 25,200 s, and a grating of 400 line mm^{-1} was used, covering ~ 5600 and $8800 \text{ \AA}$ with a resolution of $\sim 5.8 \text{ \AA}$. With the seeing of $\sim 1''.5$, all the observations were made with a series of exposures of 30 minutes each. On each night, low-resolution spectra were also obtained to cover the wavelength band near the Ly α emission.

The data were processed and analyzed using standard *IRAF* tasks. The A- and B-band absorption were corrected with the templates derived from the standard stars. The removal was only partial because some exposures were obtained with large air mass without corresponding spectra of nearby template stars. Spectra taken on different dates were combined, using exposure times as the weight. The spectra of HS 1700+64 and Q0302-003 are displayed respectively in Fig. 1 and 2. The spectrum of HS 1700+64 covers too small a wavelength region shortward of the C IV to allow an accurate determination of the continuum level. We used another spectrum taken in 1994 June with the same telescope that covers wavelengths as low as $5000 \text{ \AA}$. The fitted continuum matches that spectrum well. We therefore feel that the continuum used in the fitting is appropriate.

The color excess E_{B-V} was calculated as 0.05 for HS 1700+64 and 0.12 for Q0302-003, according to the estimated column density of Galactic neutral hydrogen (Stark et al. 1992). Such a conversion was based on the average Galactic dust-to-gas ratio derived by Bohlin, Savage, & Drake (1978), which yields a larger correction than using the formulation of Burstein & Heiles (1978). The blue part of the He II $\lambda 304$ profile is absorbed by the intergalactic He II absorption, so we only fitted the red part. The fitting results are presented in Table 1. The Ly α measurements were made using the spectra of lower resolution taken in the same observing runs.

For HS 1700+64, the fitted He II $\lambda 304$ profile has a FWHM of $\sim 12\,000 \text{ km s}^{-1}$, which matches that of the Ly α emission (FWHM $\sim 11\,500 \text{ km s}^{-1}$; Reimers et al. 1989). In the optical spectrum there is no noticeable emission feature at the expected He II $\lambda 1640$ wavelength. To obtain an upper limit, we assumed a Gaussian profile with a fixed centroid wavelength at $6138 \text{ \AA}$ and a FWHM of $12\,000 \text{ km s}^{-1}$. The He II $\lambda 1640$ flux is $(0.3 \pm 2.8) \times 10^{-15} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$, yielding an $I(\lambda 304)/I(\lambda 1640)$ ratio of greater than 20. Such a value may not be very reliable as a slight shift in the continuum level may change the ratio significantly. The current broad-band data of

standard stars may not allow a confirmation of broad features at 1% level. Therefore the result may not be inconsistent with the canonical value of 10. The emission centered around 6385 Å does not correspond to known emission features in quasars, and we tentatively linked it to Si I λ 1701 (Verner, Barthel & Tytler 1994), or a Fe II feature (Francis et al. 1991).

For Q0302-003, there is a significant narrow component for the Ly α and C IV emission. Each of the Ly α , C IV and N V was fitted with a pair of Gaussian components. The narrow component has a FWHM of $\sim 2000 \text{ km s}^{-1}$ and accounts for about a half of the line intensity. The derived $I(\lambda 304)/I(\lambda 1640)$ ratio is ~ 1.1 , significantly lower than the predicted value with pure recombination.

3. Discussions

The $I(\lambda 304)/I(\lambda 1640)$ ratio is quite different in these two quasars. Indeed the line profiles in Q0302-003 are narrower, making it easier to identify the weak He II λ 1640 feature. The S/N level is high enough that a He II λ 1640 feature should be detected even with a line width of $\sim 12000 \text{ km s}^{-1}$. In Fig. 1 the profile of an assumed He II λ 1640 feature is plotted, with an intensity 20% of the He II λ 304, which should have been detected. It appears that difference is not simply attributable to line widths.

The intensity of He II λ 304 emission is affected by the Lyman line and continuum absorption by numerous intervening absorbers along the line of sight. This can be corrected if a high-resolution spectrum at longer wavelengths yields a list of absorption lines. Our estimate, based on the statistical result of Møller & Jakobsen (1990), suggests an optical depth of 0.2 at a rest-frame wavelength of 300 Å for a $z=3.3$ quasar. Therefore, this Lyman-Valley correction is not very significant.

The UV and optical observations are not simultaneous, and both quasars are probably variable. A comparison of the UV spectra of HS 1700+64 obtained between 1991 and 1995 finds a significant discrepancy in flux level, and that between the optical spectra taken between 1994 and 1996 shows that the Ly α equivalent width varies by a factor of 2. The C IV equivalent width of Q0302-003 has increased by $\sim 50\%$ as compared with the data of Sargent, Steidel & Boksenberg (1989). Furthermore, the photometric quality of our optical spectra is questionable. A typical light loss with a small slit during an optical spectroscopic observation is $\sim 15\%$. These factors add uncertainties to the $I(\lambda 304)/I(\lambda 1640)$ ratio. The derived line ratio is also subject to the reddening formulation. If we use the formula of Burstein & Heiles (1978), this ratio would be even lower.

We have carried out photoionization calculations (Ferland 1996) with various parameters. With a broad range of the density, column density, flux and shape of the ionizing continuum, the $I(\lambda 304)/I(\lambda 1640)$ ratio varies within a narrow range between 9 and 11. It is therefore not practical to attribute the observed low value to special conditions in the line-emitting region. Note that a part of the He II λ 304 emission may receive a contribution from O III λ 305 emission that is produced

by Bowen fluorescence mechanism (Eastman & MacAlpine 1985). If this were the case, the actual $I(\lambda 304)/I(\lambda 1640)$ ratio would be even lower.

The low $I(\lambda 304)/I(\lambda 1640)$ ratio in Q0302-003 may signal internal reddening in the line-emitting region. Dust grains with dimensions of $\sim 3 \times 10^{-6}$ cm are believed to produce Galactic extinction (Savage & Mathis 1979) which generally follows a $1/\lambda$ law. If intrinsic extinction is produced by even smaller grains, and the wavelength dependence of the extinction law applies to wavelengths as short as 300 Å, then an $E_{E-V} = 2.5$, where E denotes a band around 300 Å, is needed to explain the discrepancy between the observed and theoretical ratio of $I(\lambda 304)/I(\lambda 1640)$. This would translate into $E_{B-V} = 0.5$. If this is the case, significant presence of dust may be a reality in the broad-line region of some high-redshift quasars.

The quantitative formalism should be more complicated than that. While the extinction curves between 1 μm and 1000 Å can be approximated with a $1/\lambda$ law, very little is known about the extinction properties below 1000 Å. Hawkins & Wright (1989) and Martin & Rouleau (1991) calculated the EUV extinction curve for graphite-silicate dust. Their results show *decreasing* extinction from 1000 Å to 100 Å. Pei (1992) suggested that a numerical formula can be applied to other galaxies, possibly to those of higher redshifts, without assuming a Galactic dust-to-gas ratio. If such extinction is real, the same extinction affects the intensity of other UV lines as well. The intrinsic hydrogen $\text{Ly}\alpha/\text{H}\alpha$ ratio in these objects may actually be higher than the observed ones. Likely, the average ratio would be ~ 6 , closer than the theoretical value of ~ 10 . This will, in turn, help understand the classical $\text{Ly}\alpha/\text{H}\alpha$ puzzle. If significant reddening does exist in the quasar broad-line region, the observed $\text{Ly}\alpha/\text{H}\alpha$ ratio may be corrected upward by an additional factor of ~ 2 .

Even for these two quasars, the $I(\lambda 1216)/I(\lambda 304)$ ratio is very different. In HS 1700+64, this ratio is about 3, while in Q0302-003 it is about 30. Significant extinction in the EUV band can explain both the abnormal line ratios. The $I(\lambda 1216)/I(\lambda 304)$ ratio in both objects is 30 or higher, consistent with photoionization models. Therefore, the likely cause for the low ratio in Q0302-003 is extinction in the broad line region by very small grains.

It may not be coincidental that narrow line widths and possibly significant reddening are present in the same object. Seyfert 2 galaxies often show a higher degree of extinction (Osterbrock 1989), and the narrow-line region in Seyfert-1 galaxies generally exhibit a more significant reddening effect than the broad-line region (Binette, et al. 1993). The spectrum of Q0302-003 shows a significant narrow component with $\text{FWHM} \sim 2000 \text{ km s}^{-1}$ for major emission lines. Although this line width is not considered very narrow for Seyfert galaxies, it is for high- z quasars. Generally, narrow lines in high- z quasars (Sargent, et al. 1989) are not as common as in Seyfert galaxies. For example, Wills et al. (1993) found no detection of narrow line components in their radio-loud quasars of $0.26 < z < 0.77$. They suggested a significant reddening with $E_{B-V} \simeq 0.5$. We suggest that the geometry of the line-emitting region in high- z quasars resembles that in the low-luminosity active galaxies, with the presence of dust mostly in the outer part.

Making an analogy of Seyfert galaxies and some low-redshift quasars, we suggest that Q0302-003 has a narrow-line region which contains a significant amount of dust. We suggest that the geometry of the line-emitting region in high- z quasars resembles that in the low-luminosity active galaxies, with the presence of dust mostly in the outer part (Binette, et al. 1993).

Does reddening apply to the EUV continuum? Recent studies (Netzer, et al. 1995; Bechtold 1997) found that the $\text{Ly}\alpha/\text{H}\beta$ ratio ranges between about 1 and 40 and is approximately proportional to $f(1216)/f(4861)$, the ratio of continuum flux at adjacent points (Bechtold 1997). Such a correlation may suggest a possible reddening effect that applies to both the continuum and lines. In such cases, the equivalent widths of concerned lines should be fairly constant. Given the significant difference in the equivalent widths of He II lines in our quasar samples, we see no compelling reason that the continuum emission from the central source is heavily reddened. Significant reddening would also affect the intensities of infrared lines, and future studies of these lines may provide additional evidence for fine dust in the quasar environment.

Support for this work has been provided in part by NASA contract NAS-5-27000 to the Johns Hopkins University and NASA grant AR-5284.01-93A from the Space Telescope Science Institute, which is operated by the Association of Universities of Research in Astronomy, Inc., under NASA contract NAS5-26555. We are very grateful to P. S. Smith for his assistance during the observations.

REFERENCES

- Baldwin, J. A. 1977, MNRAS, 178, 67P
- Bechtold, J., et al. 1997 in Emission Lines in Active Galaxies: New Methods and Techniques, ed. B.M. Peterson, F.-Z. Cheng, and A.S. Wilson (San Francisco: ASP), 123
- Binette, L., Wang, J., Villar-Martin, M., Martin, P. G., & Margris C. G. 1997, ApJ, 414, 535
- Bohlin, R.C., Savage, B. D., & Drake, J. F. 1978, ApJ, 224, 132
- Burstein, D., & Heiles, C. 1978, ApJ, 225, 40
- Cheng, F. Z., Danese, L., & de Zotti, G. 1983, MNRAS, 204, 13P
- Davidson, A. F., Hartig, G. F., & Fastie, W. G. 1977, Nature, 269, 203
- Davidson, A. F., Kriss, G. A., & Zheng, W. 1996, Nature, 380, 47
- Eastman, R. G., & MacAlpine, G. M. 1985, ApJ, 299, 785
- Ferland, G. J. 1996, Univ. Kentucky Dept. of Phys. & Astron. Internal Report
- Ferland, G. J., & Shields, G. A. 1985, in Astrophysics of Active Galaxies and Quasi-Stellar Objects, ed. J. S. Miller (Mill Valley, CA: University Sciences), 157
- Francis, P. J., Hewett, P. C., Foltz, C. B., Chaffee, F. H., & Weymann, R.J. 1991, ApJ, 373, 465
- Hawkins, I., & Wright, E. L. 1989, in Extreme Ultraviolet Astronomy, ed. R. F. Malina & S. Bowyer (New York: Pergamon), 333
- Hogan, C. J., Anderson, S. F., & Rugers, M. H. 1997, AJ, 113, 1495
- Jakobsen, P., Boksenberg, A., Deharveng, J. M., Greenfield, P., Jedrzejewski, R., & Paresce, F. 1994, Nature, 370, 35
- MacAlpine, G. M. 1985, in Astrophysics of Active Galaxies and Quasi-Stellar Objects, ed. J. S. Miller (Mill Valley, CA: University Sciences), 259
- Martin, P. G. & Rouleau, F. 1991, in Extreme Ultraviolet Astronomy, ed. R. F. Malina & S. Bowyer, (Oxford: Pergamon), 341
- Møller, P. & Jakobsen, P. 1990, A&A, 228, 299
- Netzer, H., Elitzur, M., & Ferland, G. J. 1985, ApJ, 299, 752
- Netzer, H., & Laor, A. 1993, ApJ, 404, L51
- Netzer, H., et al. 1995, ApJ, 448, 27

- Osterbrock, D. E. 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (Mill Valley: University Science Books)
- Pei, Y. C. 1992, *ApJ*, 395, 130
- Puetter, R. C., Smith, H. E., Soifer, B. T., Willner, S. P. & Pipher, J. L. 1978, *ApJ*, 226, L53
- Reimers, D., Köhler, S., Wisotzki, L., Groote, D., Rodriguez-Pascual, P., & Wamsteker, W. 1997, *A&A*, 327, 890
- Reimers, D., et al. 1989, *A&A*, 218, 71
- Rudy, R. J., Schmidt, G. D., Stockman, H. S., & Moore, R. L. 1983, *ApJ*, 271, 59
- Sargent, W. L. W., Steidel, C. C., & Boksenberg, A. 1989, *ApJS*, 69, 703.
- Savage, B. D. & Mathis, J. S. 1979, *ARAA*, 17, 73
- Stark, A. A., Gammie, C. F., Wilson, R. W., Bally, J., Linke, R. H., Heiles, C., & Hurwitz, M. 1992, *ApJS*, 79, 77
- Tytler, D., Fan, X.-M., Burles, S., Cotrell, L., Davis, C., Kirkman, D., & Zuo, L. 1995, in *QSO Absorption Lines*, ed. G. Meylan, (Berlin: Springer), 289
- Verner, D. A., Barthel, P. D., & Tytler, D. 1994, *A&AS*, 108, 287
- Wills, B. J., et al. 1993, *ApJ*, 410, 534

Table 1. Emission Line Intensities^a

Line	Wavelength Å	HS 1700+64		Q0302–003	
		Flux	EW	Flux	EW
He II ^b	304	63 ± 13	32	4.9 ± 1.0	8
Ly α	1216	202 ± 8	142	155 ± 9^c	305
C IV	1549	117 ± 15	116	23.5 ± 1.8^c	91
He II	1640	0.3 ± 2.8	0.3	4.5 ± 0.4	16

^aFluxes are in units of 10^{-15} ergs s^{−1} cm^{−2} Å^{−1}, corrected for Galactic absorption.

^bTotal intensity, based on fit to the red part of the profile.

^cA narrow component accounts for about a half of the intensity.

Fig. 1.— Optical spectrum of HS 1700+64 ($z=2.743$). The atmospheric B-band absorption, marked with an Earth symbol, has been partially corrected. The fitted power-law continuum is plotted in dashed curve, and the fitting windows are marked. An assumed He II $\lambda 1640$ profile, with $1/5$ of the intensity of He II $\lambda 304$ line is plotted in dotted curve at the bottom, to allow an assessment of the He II $\lambda 1640$ intensity. A tentative line identification is marked with a question mark.

Fig. 2.— Optical spectrum of Q0302-003 ($z=3.286$). The atmospheric A- and B-band absorption features, marked with Earth symbols, have been partially corrected. The fitted power-law continuum is plotted in dashed curve, and the fitting windows are marked.




Fig. 1.—




Fig. 2.—